



Nienhuis Goyens, H.J. & R. Guerrero Caballero (1985). A quantitative analysis of annual phytoplankton cycle of the Magdalena lagoon complex (Mexico). Journal of Plankton Research, 7(4): 427-441. DOI: 10.1093/plankt/7.4.427

A quantitative analysis of annual phytoplankton cycle of the Magdalena lagoon complex (Mexico)

Hedrik J. Nienhuis Goyens & Rafael Guerrero Caballero

An annual cycle of the micro- and nanophytoplankton cell densities of whole water samples in combination with the species composition, distribution and diversity of the microphytoplankton fraction in the Magdalena lagoon system, Mexico, has been studied on a bi-monthly basis. Two major patterns were detected: from November to May, high microphytoplankton densities ($0.5\text{--}1.5 \times 10^6$ cells l^{-1}) prevail throughout a large part of the investigated area; from late spring to late autumn cell densities are much lower ($5\text{--}250 \times 10^3$ cells l^{-1}). During each period a number of microplankton assemblages occur, each with its characteristic diversity and stability. The distribution and density of the microphytoplankton is clearly related to the presence of nutrient-rich water pockets and the prevailing winds for part of the year, as documented in earlier hydrological studies. The nanophytoplankton played a minor role during most of the year; this fraction dominated quantitatively in only a few cases. Based on these data, the Magdalena lagoon complex can be considered a very productive area. This is most exclusively and by phytoplankton. The quantitative existence of several important species, feeding the food web of this coastal lagoon system.

Palabras clave: Phytoplankton, Magdalena lagoon

Para obtener copia del documento contacta con el autor (ngoyens@ipn.mx) o con el personal de la biblioteca (bibliocicimar@ipn.mx).