



Poortvliet, M., **F. Galván Magaña**, G. Bernardi, D.A. Croll & J.L. Olsen (2011). Isolation and characterization of twelve microsatellite loci for the Japanese Devilray (*Mobula japonica*). Conservation Genetics Resources, 3(4): 733-735. DOI: 10.1007/s12686-011-9445-0

Isolation and characterization of twelve microsatellite loci for the Japanese Devilray (*Mobula japonica*)

Marloes Poortvliet, Felipe Galván Magaña, Giacomo Bernardi, Donald A. Croll & Jeanine L. Olsen

Twelve polymorphic microsatellites loci were characterized for *Mobula japonica* (Japanese Devilray) using an enrichment protocol. All but two loci were in Hardy–Weinberg equilibrium with no evidence of linkage disequilibrium or null-alleles for a sample of 40 individuals from two populations. The number of alleles varied from 5 to 28. Expected heterozygosity ranged from 0.2332 to 0.9589, making these microsatellite loci good candidates for population genetic studies.

Palabras clave: Elasmobranch, microsatellite, *Mobula japonica*, Population genetics, Polymorphism

Para obtener copia del documento contacta con el autor (fgalvan@ipn.mx) o con el personal de la biblioteca (bibliocicimar@ipn.mx).